VOL. 582, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY NOVEMBER 6, 1992

(Biomedical Applications, Vol. 120, Nos. 1+2)

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Research Alert, Science Citation Index and Trends in Biotechnology)

Publisher's Note V

Determination of dopamine and two of its acidic metabolites by high-performance liquid chromatography with electrochemical detection: a retrospective article (Chromatography Classic)
by O. Magnusson and L. B. Nilsson (Södertälje, Sweden) and D. Westerlund (Uppsala, Sweden) (Received May 19th, 1992) 1

Urinary testosterone measurement by gas chromatography after solid-phase extraction and high-performance liquid chromatography
by E. Venturelli, A. Manzari and A. Cavalleri (Milan, Italy), M. Benzo (Cernusco sul Naviglio, Italy) and G. Secreto and E. Marubini (Milan, Italy) (Received July 8th, 1992) 7

Rapid and simple method for the determination of urinary benzoic and phenylacetic acids and their glycine conjugates in ruminants by reversed-phase high-performance liquid chromatography
by M. J. Arín, M. T. Diez and J. A. Resines (León, Spain) (Received July 16th, 1992) 13

Simultaneous measurement of serotonin, catecholamines and their metabolites in cat and human plasma by *in vitro* microdialysis–microbore high-performance liquid chromatography with amperometric detection
by F.-C. Cheng, L.-L. Yang, F.-M. Chang, L.-G. Chia and J.-S. Kuo (Taichung, Taiwan) (Received July 21st, 1992) . 19

Microbore liquid chromatography with dual electrochemical detection for the determination of serotonin and 5-hydroxyindoleacetic acid in rat brain dialysates
by S. Sarre and Y. Michotte (Brussels, Belgium), C. A. Marvin (Lier, Belgium) and C. Ebinger (Brussels, Belgium) (Received July 29th, 1992) 29

Determination of D-alanine and D-glutamic acid in biological samples by coupled-column chromatography using β-cyclodextrin as mobile phase additive
by A. M. Rizzi, P. Briza and M. Breitenbach (Vienna, Austria) (Received July 30th, 1992) 35

Determination of free amino acid enantiomers in rat brain and serum by high-performance liquid chromatography after derivatization with N-*tert.*-butyloxycarbonyl-L-cysteine and *o*-phthaldialdehyde
by A. Hashimoto (Ibaraki, Japan) and T. Nishikawa, T. Oka, K. Takahashi and T. Hayashi (Tokyo, Japan) (Received July 15th, 1992) 41

Quantitative analysis of collagen and elastin cross-links using a single-column system
by T. J. Sims and A. J. Bailey (Bristol, UK) (Received July 18th, 1992) 49

High-performance liquid chromatographic determination of dihydroorotate dehydrogenase of *Plasmodium falciparum* and effects of antimalarials on enzyme activity
by I. Ittarat, H. K. Webster and Y. Yuthavong (Bangkok, Thailand) (Received July 8th, 1992) 57

Evaluation of the chromatographic procedure for the preparation of a high-purity C1-esterase inhibitor concentrate from cryosupernatant plasma
by L.-C. Teh and M. Froger (Auckland, New Zealand) (Received July 15th, 1992) 65

Rapid purification yielding highly active 17β-hydroxysteroid dehydrogenase: application of hydrophobic interaction and affinity fast protein liquid chromatography
by F. Yang, D.-W. Zhu, J.-Y. Wang and S.-X. Lin (Ste-Foy, Canada) (Received July 12th, 1992) 71

Separation of serum bilirubin species by micellar electrokinetic chromatography with direct sample injection
by N. Wu, T. Wang, R. A. Hartwick and C. W. Huie (Binghamton, NY, USA) (Received July 15th, 1992) 77

Polypeptide marker and disease patterns found while mapping proteins in ascitis
by A. Toussi, N. Paquet, O. Huber and S. Frutiger (Geneva, Switzerland), J.-D. Tissot (Lausanne, Switzerland) and G. J. Hughes and D. F. Hochstrasser (Geneva, Switzerland) (Received July 15th, 1992) 87

(Continued overleaf)

Determination and identification of amiloride in human urine by high-performance liquid chromatography and gas chromatography–mass spectrometry
by H. Bi, S. F. Cooper and M. G. Côté (Pointe-Claire, Canada) (Received July 27th, 1992) 93

Determination of drugs from urine by on-line immunoaffinity chromatography–high-performance liquid chromatography–mass spectrometry
by G. S. Rule and J. D. Henion (Ithaca, NY, USA) (Received July 17th, 1992) 103

Determination of chloramphenicol in muscle using a particle beam interface for combining liquid chromatography with negative-ion chemical ionization mass spectrometry
by B. Delépine and P. Sanders (Fougeres, France) (Received August 11th, 1992) 113

On-line dialysis and weak cation-exchange enrichment of dialysate. Automated high-performance liquid chromatography of pholcodine in human plasma and whole blood
by A. T. Andresen, M. Krogh and K. E. Rasmussen (Oslo, Norway) (Received August 12th, 1992) 123

High-performance liquid chromatographic determination of cyanide in human red blood cells by pre-column fluorescence derivatization
by A. Sano, N. Takimoto and S. Takitani (Tokyo, Japan) (Received August 11th, 1992) 131

Rapid and sensitive determination of coumarin and 7-hydroxycoumarin and its glucuronide conjugate in urine and plasma by high-performance liquid chromatography
by D. A. Egan and R. O'Kennedy (Dublin, Ireland) (Received August 11th, 1992) 137

High-performance liquid chromatographic determination of nicotinamide and its metabolites in human and murine plasma and urine
by M. R. L. Stratford and M. F. Dennis (Northwood, UK) (Received August 5th, 1992) 145

Isolation and structural elucidation of the geometrical isomers of lutein and zeaxanthin in extracts from human plasma
by F. Khachik (Beltsville, MD and Washington, DC, USA), G. Englert (Basel, Switzerland) and C. E. Daitch, G. R. Beecher, L. H. Tonucci and W. R. Lusby (Beltsville, MD, USA) (Received July 13th, 1992) 153

Determination of cocaethylene, cocaine and their metabolites in rat serum microsamples by high-performance liquid chromatography, and its application to pharmacokinetic studies in rodents
by C. E. Lau (New Brunswick, NJ, USA) (Received July 13th, 1992) 167

High-performance liquid chromatographic analysis of methocarbamol enantiomers in biological fluids
by S. Alessi-Severini, R. T. Coutts, F. Jamali and F. M. Pasutto (Edmonton, Canada) (Received July 22nd, 1992) . . 173

Determination of acetazolamine in human urine samples by reversed-phase high-performance liquid chromatography in the presence of xanthines
by R. Herráez-Hernández, P. Campins-Falcó and A. Sevillano-Cabeza (Valencia, Spain) (Received August 11th, 1992) 181

High-performance liquid chromatographic determination of diuretics in urine by micellar liquid chromatography
by E. Bonet Domingo, M. J. Medina Hernández, G. Ramis Ramos and M. C. García Alvarez-Coque (Valencia, Spain) (Received August 6th, 1992) 189

Quantitation of the novel anticonvulsant remacemide in rat and dog plasma and urine: application of the plasma methodology to measure the plasma protein binding of remacemide
by T. C. M. Wilson, M. S. Eisman and G. E. Machulskis (Rochester, NY, USA) (Received August 11th, 1992) . . . 195

Simple and sensitive high-performance liquid chromatographic method for the determination of diltiazem and six of its metabolites in human plasma
by M. D. Hussain, Y. K. Tam, B. A. Finegan and R. T. Coutts (Edmonton, Canada) (Received August 3rd, 1992) . . 203

Determination of thalidomide in plasma and blood by high-performance liquid chromatography: avoiding hydrolytic degradation
by T. Eriksson, S. Björkman, Å. Fyge and H. Ekberg (Malmö, Sweden) (Received June 16th, 1992) 211

High-performance liquid chromatographic method for quantitating plasma levels of amiloride and its analogues
by M. A. Alliegro, K. D. Dyer, E. J. Cragoe, Jr., B. M. Glaser and M. C. Alliegro (Baltimore, MD, USA) (Received July 15th, 1992) 217

High-performance liquid chromatographic analysis of the anticancer drug oxantrazole in rat whole blood and tissues
by E. E. Hassan and J. M. Gallo (Athens, GA, USA) (Received July 10th, 1992) 225

Short Communications

Improved sample preparation for the testosterone hydroxylation assay using disposable extraction columns
by F. Oesch, H. Wagner, K. L. Platt and M. Arand (Mainz, Germany) (Received July 3rd, 1992) 232

Determination of plasma homovanillic acid by liquid chromatography with electrochemical detection
by R. N. Gupta and C. Welton (Hamilton, Canada) (Received July 23rd, 1992) 236

Ion-pair high-performance liquid chromatography of diaminopimelic acid in hydrolysates of physiological samples
by M. E. R. Dugan, W. C. Sauer, K. A. Lien and T. W. Fenton (Edmonton, Canda) (Received August 12th, 1992) . 242

Rapid and simple method for the determination of α_1-acid glycoprotein in serum by column liquid chromatography
by S. Kishino, Z. S. Di, M. Sugarawa, K. Iseki, K. Miyazaki, S. Kakinoki, A. Nomura and A. Kitabatake (Sapporo, Japan) (Received July 17th, 1992) 246

Improved procedure for the drying and storage of polyacrylamide slab gels
by Ľ. Škultéty and R. Toman (Bratislava, Czechoslovakia) (Received July 31st, 1992) 249

High-performance liquid chromatographic monitoring of metabolic products resulting from the treatment of mouse hepatoma cells with N^6-cycloalkylated nucleosides
by R. Thedford, L. P. Fauntleroy, Y. L. Murray, M. Stokes and C. L. Thornton (Atlanta, GA, USA) (Received August 5th, 1992) 253

Simple high-performance liquid chromatographic method for the determination of captopril in biological fluids
by G. Shen, T. Weirong and W. Shixiang (Shanghai, China) (Received August 11th, 1992) 258

Improved high-performance liquid chromatographic determination of ciprofloxacin and its metabolites in human specimens
by G. Mack (Westmead, Australia) (Received July 27th, 1992) 263

Automated high-performance liquid chromatographic method for the determination of mianserin in plasma using electrochemical detection
by L. W. Brown, H. K. L. Hundt and K. J. Swart (Bloemfontein, South Africa) (Received July 22nd, 1992) 268

Determination of Taxotere in human plasma by a semi-automated high-performance liquid chromatographic method
by J. C. Vergniol, R. Bruno, G. Montay and A. Frydman (Antony, France) (Received July 8th, 1992) 273

Related articles published in Journal of Chromatography, Vols. 623 and 624 279

Author Index 281

Compound Index 284